



DANGEROUS POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



LAST SUPPER SUPREME 50

ZINC Phosphide (Broadacre) Mouse Bait Pellets

ACTIVE CONSTITUENT: ZINC PHOSPHIDE

For the control of heavy infestations of mice in agricultural situations.

THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROL FOR SCHEDULE 7 POISONS.

NET CONTENTS: 125 kg 500 kg

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover).
DO NOT apply bait in a trail.
DO NOT apply bait if heavy rain is imminent.
DO NOT apply to the outer 50 m of crop or within 50 m of native vegetation. A 50 m buffer zone in crop must be employed to reduce risk to birds and other non-target animals.
DO NOT apply bait unless monitoring of mouse numbers indicates the potential for crop damage of economic significance.

SITUATION	PEST	RATE	CRITICAL COMMENTS
Grain crops, legume crops, canola, safflower, nut crops and pasture. (Note: for all situations - only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seeds have been sown into crop stubble or other similar ground cover)	Mice	1 kg/ha to achieve an even coverage of 2 pellets /3m ²	Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2 pellets/ 3 m2. It is imperative that all attempts are made to apply bait as evenly as possible to maximise product efficacy and minimise risks to non-target animals. This product will provide a high level of control in most situations. However, in areas of extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Re-treatment after 14 days may therefore be required in certain circumstances. It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying the bait and that post-baiting estimates are conducted to assess the effectiveness of the control operation. For information on the most effective methods for monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control. Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.

GENERAL INSTRUCTIONS

Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS, THE ENVIRONMENT AND LIVESTOCK

Do not contaminate dams, rivers, streams or waterways with product or the used container.

STORAGE AND DISPOSAL

The contents of each container should be used within 3 months of opening. Store in the closed original container in a dry cool well-ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any sources of heat or ignition. Dispose of used containers by crushing and burying below one metre in a disposal pit specifically marked for this purpose, clear of waterways, vegetation and roots. Excess or unused baits must be buried below one metre. Empty containers and product must not be burnt.

SAFETY DIRECTIONS

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames – forms toxic gas. When opening the container and using the product wear elbow length PVC gloves and a full face respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each days use wash gloves and respirator and if rubber wash with detergent and warm water.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. Do not give direct mouth to mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy viva or one-way mask. Resuscitate in a well ventilated area.

SAFETY DATA SHEET

Additional information is listed in the SDS which can be contained by the supplier.

CONDITIONS OF SALE

The use of Last Supper Supreme 5- Zinc Phosphide (Broadacre) Mouse Bait Pellets being beyond the control of the manufacturer, no warranty expressed or implied is given by Wilhelm Rural Pty Ltd, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wilhelm Rural Pty Ltd accepts no responsibility for any consequence whatsoever from the use of this product.

Manufactured & Distributed by

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN. DO NOT USE THIS PRODUCT IN THE HOME GARDEN.

WITHHOLDING PERIOD

DO NOT HARVEST ANY CROPS OR ALLOW LIVESTOCK TO GRAZE BAITED AREAS FOR 14 DAYS AFTER APPLICATION.



BN: _____ DOM: _____